

A possible anisotropy in blackbody radiation viewed through non-uniform gaseous matter

T K Rai Dastidar*

Indian Association for the Cultivation of Science, Calcutta 700032, India

Abstract

A non-local gauge symmetry of a complex scalar field, which can be trivially extended to spinor fields, was demonstrated in a recent paper (Mod. Phys. Lett. **A13**, 1265 (1998) ; hep-th/9902020). The corresponding covariant Lagrangian density yielded a new, non-local Quantum Electrodynamics. In this Letter we show that as a consequence of this new QED, a blackbody radiation viewed through gaseous matter appears to show a slight deviation from the Planck formula, and propose an experimental test to check this effect. We also show that a non-uniformity in this gaseous matter distribution leads to an (apparent) spatial anisotropy of the blackbody radiation.

* Electronic address : mstkrd@mahendra.iacs.res.in

I. Introduction

We have recently [1] shown how a non-local model of Quantum Electrodynamics (QED), whose phenomenological form had been given earlier [2], results under a generalization of the usual U(1) gauge transformation to a non-local one :

$$\exp(-ie\Lambda(x)) \rightarrow \exp(-ie\Lambda|x\rangle\langle x'|). \quad (1)$$

where $x \equiv (\mathbf{r}, t)$, $x' \equiv (\mathbf{r}', t')$. It was shown in [2] that a covariant derivative $\mathcal{D}_\mu = \partial_\mu + d_\mu + ie\mathcal{A}_\mu|x\rangle\langle x'|$ (where $d_\mu \equiv \frac{\partial}{\partial x'^\mu}$) can be used to write down a covariant Lagrangian for a complex scalar field $\phi(x)$:

$$\mathcal{L} = -\frac{1}{4}\mathcal{F}_{\mu\nu}\mathcal{F}^{\mu\nu} + \mathcal{D}_\mu\phi^*\mathcal{D}^\mu\phi \quad (2)$$

where

$$\mathcal{F}^{\mu\nu} \equiv \mathcal{F}^{\mu\nu}|x\rangle\langle x'| = (\partial^\mu + d^\mu)\mathcal{A}^\nu|x\rangle\langle x'| - (\partial^\nu + d^\nu)\mathcal{A}^\mu|x\rangle\langle x'| \quad (3)$$

is the non-local electromagnetic field tensor. Under the gauge transformation

$$\mathcal{A}_\mu|x\rangle\langle x'| \rightarrow \mathcal{A}_\mu|x\rangle\langle x'| + \partial_\mu\Lambda|x\rangle\langle x'| + d_\mu\Lambda|x\rangle\langle x'|. \quad (4)$$

$\mathcal{F}^{\mu\nu}$ remains gauge invariant if and only if

$$[\partial^\mu, d^\nu] = 0. \quad (5)$$

The relation (5) may therefore be considered as the necessary and sufficient condition for the non-local gauge transformation (1) to be a symmetry of the field. This relation (5), which implies that the two space-time points x, x' are completely independent (i.e. need not lie within the light cone of one another), shows that the non-local correlation must be of the Einstein-Podolsky-Rosen (EPR)-type. It was shown in [1] that :

1. A correlated two-photon absorption *linear in intensity* emerges as a consequence of this non-local symmetry of the radiation field. As mentioned in [1] (where references of related work in ordinary QED have been given), experimental evidence of two-photon absorption of such kind has been obtained recently [3].

2. A non-thermodynamic Time's Arrow in the quantum (atomic) level emerges as a necessary condition for energy conservation in such correlated two-photon absorption processes, thus relegating the principle of causality to a corollary of the principle of energy conservation.
3. The requirement for the EPR-type nonlocal correlation, which is in agreement with Bell's theorem, is also a necessary condition for energy conservation in interaction of matter with radiation involving a correlated two-photon exchange.

In this Letter it is shown that if we look, through gaseous matter, at a blackbody radiation field (with an arbitrary temperature T), then — as a consequence of this correlated two-photon absorption by the gas — the spectrum of the radiation would appear to us to be slightly deviated from the Planck formula, the amount of the deviation being directly proportional to the density of the gaseous matter. Furthermore, a non-uniform distribution of this matter would appear to us as an apparent spatial anisotropy of this radiation field temperature.

II. Theory

The energy density of blackbody radiation at temperature T (deg K) within a frequency band $\nu \rightarrow \nu + d\nu$ is given by the Planck formula

$$E(\nu)d\nu = A \left(\frac{u^3}{e^u - 1} \right) d\nu, \quad \text{where} \quad A = \frac{8\pi}{h^2} \left(\frac{kT}{c} \right)^3, \quad u = \frac{h\nu}{kT}. \quad (6)$$

The function $u^3/(e^u - 1)$ is maximum at $u \simeq 2.821\,439$, i.e. the blackbody temperature T is related to the frequency ν_m at the peak by $T \simeq h\nu_m/(2.821\,439k)$.

We recall from [1] that when matter-radiation interaction satisfies the condition of a second-order coherence, the symmetry of invariance under a non-local gauge transformation becomes prominent and causes two-photon absorption/s linear in the intensity. As a result, when a blackbody radiation reaches an observer through a gaseous absorber under circumstances (see discussion in section III) that the above condition is fulfilled, a coherent absorption of two photons of frequency $\frac{1}{2}\nu$ would be superposed as an absorption structure

at a frequency ν upon the blackbody spectrum as viewed by the observer. Under the simplest assumption that the absorption coefficient z is the same over the entire spectral range of interest, the loss $F(\nu)$ as a consequence of the correlated absorption of two photons each of frequency $\frac{1}{2}\nu$ would be given by

$$F(\nu) = zE(\nu/2) = zA \left(\frac{y^3}{e^y - 1} \right) \quad (7)$$

where $y = \frac{1}{2}u$. As a result the observed blackbody radiation density reduces to

$$G(\nu) = E(\nu) - F(\nu) = Af(u, z) \quad (8)$$

where

$$f(u, z) = \frac{u^3}{e^u - 1} - \frac{\frac{1}{8}u^3}{e^{\frac{1}{2}u} - 1}z \quad (9)$$

III. Results and discussion

Fig. 1 shows the functions $E(\nu)$, $F(\nu)$ and $G(\nu)$ (times $1/A$) at a temperature $T=2.728$ deg K plotted against ν , with an arbitrarily chosen absorption coefficient $z = 5\%$. It is evident that the peak of the $G(\nu)$ -curve, which represents the intensity spectrum viewed by the observer through the gaseous matter, is shifted slightly towards the left of the peak of the Planck curve $E(\nu)$, and this shift would appear to the observer as *an apparent shift ΔT in the radiation temperature* — in this particular example the temperature shift turns out to be ~ -0.039 deg K, i.e. about 1.4%. Sample calculations with different values of z show that the temperature shift $\Delta T/T \sim 0.3z$ always.

It remains to ascertain exactly under what circumstances the correlated two-photon absorption by the intervening gaseous matter can take place from the radiation field, so that the above effect can be observed. As shown in [1] (see §2 therein), the basic criterion is that the radiation field (either as generated, or as detected by the atomic electrons) should enjoy a large second-order coherence, i.e. it should be possible for two photons to be absorbed by the atomic electrons within a vanishingly small phase difference $\Delta\phi = \omega\delta t$,

where δt is the time interval between the two photon-absorption events. Essentially, this translates in practice to the condition that the photon density be significantly higher than the electron density (the latter remaining, at the same time, high enough to make the photon absorption process viable).

A little reflection shows that this density criterion can be fulfilled under three circumstances: *first* — in a laboratory setting — when an atom is placed in a radiation field so intense that the photon density is substantially higher than the electron density ; *second* — also a laboratory setting — when an atom is placed in a phase-squeezed light containing phase-correlated *signal* and *idler* photons ; and *third*, in a cosmological setting where the overall photon density is nine to ten orders of magnitude higher than the baryon or lepton density and hence, in the millimetre-wave region as considered in the above example, chances of two photon absorption events within vanishingly small $\omega\delta t$ are high. For example, if two millimetre-wave photons are absorbed within an interval of $\sim 0.1 - 1$ ps, then the phase gap between them is $\ll 1$. The first and second laboratory settings required for the second-order phase coherence and some of their experimental consequences have already been discussed in [1] ; the difference between the laboratory settings and the cosmological setting is that, whereas in the former the spatial distance between the two distinct photon-absorption events is of the order of atomic/molecular dimensions, in the latter the said distance can be of macroscopic scales.

Our exploratory calculations above have assumed that z is a constant. Even without going into the detailed dynamics needed to calculate z for any specific system, it is obvious that z depends on the density of the absorbing gas. This allows for a very simple test for the effect of an apparent shift in the observed blackbody radiation temperature as predicted above. One simply needs a hollow cavity “black body” maintained at any arbitrary temperature, with a spectrophotometer mounted inside and filled with any desired gas, and record the spectrum at different pressures of the gas. So long as the pressure is sufficiently low so that the electron density is significantly lower than the photon density, the abovementioned second-order coherence condition applies, and the *measured* blackbody temperature should show a variation with the gas pressure as given in eqn.(8). Furthermore, this pattern of linear variation of ΔT with z can be expected to hold only so long

as the atomic/molecular density of the gaseous matter remains much less than the photon density of the radiation field.

Turning now to the cosmological setting, if an observer looks at the cosmic microwave background radiation (CMBR) from within a relatively dense gaseous atmosphere, then, as a consequence of this correlated two-photon absorption (from the CMBR) by atmospheric gases, he would find a relatively large deviation from the Planck curve, so long as the atomic/molecular density remains much less than the photon density ; however, viewing from *outside* the atmosphere, no such large deviation would appear in his measurements. This seems to be corroborated by the difference between the balloon-borne measurements of the cosmic microwave background spectrum carried out by Woody and Richards [4] — note in Fig.2 therein the large deviation from the Planck curve at *twice* the peak frequency — and the measurements recorded by the *COBE* (Cosmic Background Explorer) satellite [5]. Of course, in the absence of any specific experimental data of the type mentioned in the earlier paragraph, no precise reason for the difference between the two sets of measurements can be given as yet, but we can make the following speculations. The balloon measurements in [4] were carried out at a float pressure of ~ 2 mbar, i.e. at a height of about 120 km above sea-level, well up in the ionosphere. The peak in the CMBR spectrum in [4] is around 6 cm^{-1} , at a brightness of $\sim 10^{-11} \text{ W/cm}^2$, which corresponds to a photon flux $\sim 10^{11} \text{ cm}^{-2} \text{ sec}^{-1}$. (Dividing by c gives a photon density $\sim 3 \text{ cm}^{-3}$.) *Twice of this photon energy* can correspond to any of the following :

- i) spacing between high-lying Rydberg levels (such as $ns^2 - n'p^2, np^2 - n'd^2 \dots$) of Helium [6], where n and n' are of order ~ 40 and can be equal ;
- ii) spacing between high-lying levels of He^+ ;
- iii) spacing between rotational sublevels of high vibrational levels of electronically excited states of H_2 or H_2^+ .

Assuming an electron temperature of $\sim 1.5 \times 10^4 \text{ deg K}$ [7] and a Helium concentration of around 5 ppm in dry air, use of the Saha formula [8] for excited-state population of He and He^+ gases at the stated pressure in local thermal equilibrium yields a density of order $\sim 10^{-4} - 10^{-5} \text{ cm}^{-3}$; this is several orders of magnitude lower than the CMBR

photon density. It is difficult to estimate the corresponding figure/s for the Hydrogen molecules, in view of the uncertainty in the concentration of water vapour in the ionosphere. Nevertheless, it seems safe to assume that the population density of the rovibrationally excited levels would be much smaller than the relevant photon density. We can therefore say that the energy levels of excited Helium and hydrogen and/or their ions as present in the upper atmosphere (at ≥ 100 km altitude) *allow a coherent two-photon absorption from the CMBR, giving rise to an apparent shift in the spectrum* as given in Fig. 1 ; experimental investigations as spelt out earlier in the paper would be required to make a quantitative comparison with the data in [4] possible. (Note that we do not expect any noticeable effect in *ground-based observations*, because of the lack of ionized and/or excited species.)

Furthermore, if the density of any gaseous matter intervening between the observer and the radiation be non-uniform in space, i.e. different along different directions with respect to the observer, then this apparent shift in the blackbody radiation temperature would also vary along different directions — in other words, a non-uniformity in the absorbing matter distribution would lead to an *apparent spatial anisotropy* in the observed temperature. It is to be noted that this effect is independent of the Doppler anisotropy caused by the observer's motion through the radiation field (e.g. [9]). Of course, our theory does not cover neutrinos which are supposed to make up the *dark matter* of the universe, and hence cannot be used to fully account for the spatial anisotropies in the CMBR ; work is in progress to include the electroweak interactions within our model.

In another forthcoming work, we intend to show how, in principle, it is possible for the cosmic background radiation to impart the atomic-level time's arrow (as derived in [1] and mentioned towards the beginning of this paper) to all particles in the universe that interact with the electromagnetic field.

The author is indebted to Dr Krishna Rai Dastidar for fruitful discussions and for critically reading the manuscript.

REFERENCES

1. T K Rai Dastidar and Krishna Rai Dastidar, Mod. Phys. Letts. **A13**, 1265 (1998) ; *Errata* **A13**, 2247 (1998). Also available in the internet as : xxx.lanl.gov/abs/hep-th/9902020.
2. T K Rai Dastidar and Krishna Rai Dastidar, Abstracts of XI Int. Conf. At. Phys. (Paris, 1988), p. I-23 ; Krishna Rai Dastidar *in* “Advances in Atomic and Molecular Physics”, ed. M S Z Chaghtai (Today’s and Tomorrow’s Publishers, New Delhi, 1992) p. 49.
3. N P Georgiades, E S Polzik, K Edamatsu, H J Kimble and A S Parkins, Phys. Rev. Lett. **75**, 3426 (1995)
4. D P Woody and P L Richards, Phys. Rev. Lett. **42**, 925 (1979)
5. J C Mather et al, Astrophys. J. **420**, 439 (1994)
6. T K Rai Dastidar, unpublished calculations.
7. See e.g. *Astrophysical Formulae* by K R Lang (Springer, 1978), Table 9.
8. M N Saha, Proc. Roy. Soc. Lond. **A99**, 135 (1921)
9. P J E Peebles and D T Wilkinson, Phys. Rev. **174**, 2168 (1968)

Figure caption

Fig. 1. The Planck function $E(\nu)$ (full line), coherent two-photon absorption loss $F(\nu)$ (dashed line) with an absorption coefficient $z = 5\%$, and the resultant intensity function $G(\nu)$ (dotted line) plotted against the frequency ν . (T=2.728 deg K.) All the functions have been scaled down by the factor A (eq.(6)) to yield dimensionless quantities.

